

## Cold-Ion Spectroscopy

International Edition: DOI: 10.1002/anie.201805436 German Edition: DOI: 10.1002/ange.201805436

# **Ground-State Structure of the Proton-Bound Formate Dimer by Cold-Ion Infrared Action Spectroscopy**

Daniel A. Thomas, Mateusz Marianski, Eike Mucha, Gerard Meijer, Mark A. Johnson,\* and Gert von Helden\*

*The proton-bound* Abstract: dicarboxylate motif.  $RCOO^{-} \cdot H^+ \cdot OOCR$ , is a prevalent chemical configuration found in many condensed-phase systems. The proton-bound formate dimer  $HCOO^{-} \cdot H^{+} \cdot OOCH$  was studied utilizing cold-ion IR action spectroscopy in the range  $400-1800 \text{ cm}^{-1}$ . The spectrum obtained at ca. 0.4 K of ions captured in He nanodroplets was compared to that measured at ca. 10 K by photodissociation of Ar-ion complexes. Similar band patterns are obtained by the two techniques that are consistent with calculations for a  $C_2$  symmetry structure with a proton shared equally between the two formate moieties. Isotopic substitution experiments point to the nominal parallel stretch of the bridging proton appearing as a sharp, dominant feature near 600 cm<sup>-1</sup>. Multidimensional anharmonic calculations reveal that the bridging proton motion is strongly coupled to the flanking  $-COO^{-}$  framework, an effect that is in line with the expected change in -C=O bond rehybridization upon protonation.

he linkage formed when two neutral molecules or two anions are bound to a shared proton is a ubiquitous soft binding motif in condensed-phase chemistry. Here we are concerned with the latter case, the simplest example of which is the classic FHF- anion featuring a three-center, twoelectron bond,<sup>[1]</sup> which accommodates the bridging proton at the midpoint of the heavy atoms. A more chemically significant anionic system involves the conjugate bases of carboxylic acids, RCOO<sup>-</sup>·H<sup>+</sup>·<sup>-</sup>OOCR. These complexes are commonly observed, for example, in the anhydrous deprotonation of acidic protons in ionic liquids.<sup>[2]</sup> X-ray structures indicate that the two carboxylate groups binding the extra proton are equivalent, suggesting that, like FHF<sup>-</sup>, the proton resides at the midway point between two oxygen atoms.<sup>[3]</sup> Similarly, a recent report detailed a protein crystal structure exhibiting equivalent C–O bond lengths in a proton-bound dicarboxylate motif, potentially indicating an equally shared proton.<sup>[4]</sup> Because X-ray analysis does not reveal the location

[\*] Dr. D. A. Thomas, Dr. M. Marianski, E. Mucha, Prof. Dr. G. Meijer, Prof. Dr. G. von Helden
Fritz-Haber-Institut der Max-Planck-Gesellschaft
Faradayweg 4–6, 14195 Berlin (Germany)
E-mail: helden@fhi-berlin.mpg.de
Prof. Dr. M. A. Johnson
Sterling Chemistry Laboratory, Yale University
225 Prospect Street, New Haven, CT 06520 (USA)
E-mail: mark.johnson@yale.edu
Supporting information and the ORCID identification number(s) for
the author(s) of this article can be found under:

https://doi.org/10.1002/anie.201805436.

of the bridging proton, here we address its vibrational signature as well as those of the C–O groups residing on the flanking anions to better understand the local mechanics of this linkage. We specifically focus on the isolated HCOO<sup>-</sup>·H<sup>+</sup>·<sup>-</sup>OOCH anionic complex (hereafter denoted AHA<sup>-</sup>) and analyze its vibrational spectrum obtained using two cryogenic ion spectroscopy schemes, one involving capture and infrared photoexcitation in He droplets at 0.4 K<sup>[5]</sup> and the other based on photodissociation of the ions complexed with an Ar atom.<sup>[6]</sup>

An interesting aspect of the AHA<sup>-</sup> system is that proton attachment to one of the oxygen atoms in the carboxylate head group breaks the symmetry between the two equivalent C–O groups in the carboxylate. This in turn suggests that proton accommodation by the pair of molecular anions requires an intramolecular structural deformation. Thus, in contrast to spectroscopic characterization of the protonbound neutral complexes, AH<sup>+</sup>A, which have relied exclusively on the character of the vibrations associated with the bridging proton,<sup>[7]</sup> the carboxylates offer the opportunity to follow the evolution of the  $CO_2^-$  normal modes as the two  $CO_2^{-}$  groups deform to accommodate the bridging proton. This strategy has previously been applied to show that the intramolecular H-bond adopted by deprotonated dodecanedioic acid is, in fact, asymmetrical (that is, with the proton closer to one oxygen atom at the vibrational zero-point level).<sup>[8]</sup>

To anticipate the situation in the AHA<sup>-</sup> system, Figure 1 presents three calculated low-energy structures, which illustrate the balance of forces that drive its structure. The lowest energy form 2, for example, features a largely electrostatic binding motif in which both the OH and CH groups orient themselves towards the negatively charged oxygen atoms on the carboxylate group, reminiscent of the double contact interaction reported earlier for the I-HCO<sub>2</sub>H complex.<sup>[14]</sup> The other two isomers, on the other hand, feature a bridging O-H-O motif in which the extra proton binds to one oxygen atom on each molecular anion. These structures differ according to the orientation of the organic scaffolds, with the higher-energy conformer 3 exhibiting opposing electric dipoles, and the lower-energy conformer 1 adopting an orientation in which the formate molecular dipoles are more favorably aligned. The delicate balance of forces underlying these structures is evident when one considers that the energy ordering of these isomers changes upon inclusion of the (harmonic) vibrational zero-point energies (ZPEs). Specifically, inclusion of harmonic ZPE corrections to the Born-Oppenheimer minima brings the more symmetrical, O-H-O isomer 1 below the double-contact structure



**Figure 1.** Relative energies, zero-point energy (ZPE) corrections, and structures of the three low-energy conformations of the protonated formate dimer,  $HCOO^-$ ·H<sup>+</sup>·<sup>-</sup>OOCH. Energies are derived from complete basis set (CBS) extrapolation<sup>[9]</sup> from aug-cc-pVnZ (n=3,4,5)<sup>[10]</sup> at the MP2 level of theory and corrected for the difference between MP2<sup>[11]</sup> and CCSD(T)<sup>[12]</sup> correlation energy in the aug-cc-pVTZ basis set.<sup>[13]</sup> After inclusion of harmonic ZPE corrections, conformer **1** is predicted to be the most stable structure.

**2**. Because these systems are widely recognized to be strongly anharmonic,<sup>[7b,f,17]</sup> we have undertaken the present experimental study to establish the structure adopted at 0.4 K in He droplets by analysis of its vibrational spectrum together with multidimensional anharmonic calculations. The temperature sensitivity of these conclusions is evaluated by extending the study to the warmer conditions afforded by the Ar messenger tagging technique.

Figure 2 a,b shows the vibrational spectrum of AHA<sup>-</sup> collected between 400 and 1800 cm<sup>-1</sup> utilizing the Ar tagging and helium nanodroplet techniques, respectively. In the spectrum obtained in helium nanodroplets, an intense spectral line is observed at 605 cm<sup>-1</sup>, with other strong bands at 841, 1037, 1192, and 1684 cm<sup>-1</sup>. The Ar tagging spectrum largely agrees with that measured in helium nanodroplets, with only minor shifts in frequency observed (Supporting Information, Table S1). The relative intensities of weaker bands appear systematically reduced in the helium nanodroplet spectrum. This difference likely results from the multiple-photon absorption process utilized to generate the action signal, which is expected to result in a nonlinear dependence of signal intensity on the transition strength for a given laser fluence.<sup>[18]</sup>

As noted in the opening paragraphs, a straightforward differentiation of candidate AHA<sup>-</sup> structures exhibiting either an equally shared (1 and 3) or localized proton (2) is given by the energies of the CO<sub>2</sub><sup>-</sup> fundamentals. We focus herein on the region of the carboxylate antisymmetric stretch,  $v_{as}(COO^-)$ , and the carboxylic acid C=O stretch, v(C=O). The red and blue lines in Figure 2 denote the experimental frequencies from the literature for formate  $v_{as}(COO^-)$  and formic acid v(C=O),<sup>[15,16]</sup> which serve as limiting cases for these features in the AHA<sup>-</sup> spectrum. If the proton in AHA<sup>-</sup> is localized on an oxygen atom, C=O stretching bands are expected to occur near  $v_{as}(COO^-)$  and v(C=O), as predicted in the theoretical harmonic IR spectrum for the asymmetric conformer 2 (bands h<sub>1</sub> and h<sub>2</sub>, Figure 3 d). However, in the experimental spectrum collected in helium nanodroplets, only



*Figure 2.* Vibrational spectra of the proton-bound dimer of formate, AHA<sup>-</sup>, and its isotopically substituted variants by Ar predissociation spectroscopy and He nanodroplet ion spectroscopy. The blue and red lines mark reported experimental values for formate  $v_{as}(COO^{-})^{[15]}$  and formic acid v(C=O),<sup>[16]</sup> respectively. Strong agreement is observed between Ar tagging and He nanodroplet spectra for AHA<sup>-</sup> (a, b) and the fully deuterated analogue, ADA<sup>-</sup> (e, d). In the helium nanodroplet spectra, little change in the spectrum is observed upon deuterium substitution of the formyl hydrogen atoms (c). In contrast, a demonstrable shift of the most intense transition from 605 cm<sup>-1</sup> (b<sub>1</sub>) to 433 cm<sup>-1</sup> (d<sub>1</sub>) is observed upon substitution of the shared proton. Features labeled with \* in (d) result from proton back-exchange during ion trapping.

#### 10616 www.angewandte.org

© 2018 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim



**Figure 3.** Experimental IR spectra of AHA<sup>-</sup> (a) and ADA<sup>-</sup> (e) compared to theoretical spectra. Spectra predicted in the harmonic approximation for conformers **1** (b, f) and **2** (d, h) at the MP2/def2-TZVPP level of theory (scaling factor 0.952) fail to replicate the intense features below 1200 cm<sup>-1</sup> for both AHA<sup>-</sup> and ADA<sup>-</sup>. Theoretical spectra derived from a 4D PES for conformer **1** (c, g) show improved agreement with the experimental spectrum for both isotopologues, demonstrating the strong mode coupling in this system.

a single feature is observed with maximum intensity at 1684 cm<sup>-1</sup> (band b<sub>5</sub>), close to the midpoint of the two limiting values expected for an asymmetric geometry. A broader band is found in the spectrum obtained by Ar predissociation spectroscopy (band a<sub>5</sub>), however the prominent maxima are separated by less than  $20 \text{ cm}^{-1}$ , whereas separation of about 100 cm<sup>-1</sup> would be expected in the case of asymmetric proton localization.<sup>[8,19]</sup> Furthermore, fundamental harmonic transitions calculated for structure 1 predict two closely spaced bands in this region corresponding to in-phase and out-ofphase stretching of the two C=O bonds (band  $f_3$ , Figure 3b), features that are possibly unresolved in the helium nanodroplet spectrum. Taken together, the CO2- normal modes observed experimentally are consistent with the symmetric structure 1 for AHA<sup>-</sup> in which the proton is shared equally between the carboxylate groups.

Further support for an equally shared proton is found in the intense features measured below 1200 cm<sup>-1</sup>, which are likely associated with the vibrational modes of the shared proton.<sup>[7a,b,17,20]</sup> However, assignment of the observed bands is not straightforward, as the highly anharmonic nature of the shared proton potential precludes harmonic vibrational analysis even for one-dimensional motion, and significant coupling between normal modes is also expected.<sup>[7b, 20b, 21]</sup> Indeed, the theoretical harmonic IR spectra for conformers 1 and 2 fail to reproduce the low-frequency modes measured experimentally (Figures 3 b,d). To assist in the assignments of the features in the lower energy range, we acquired the spectra of isotopically substituted variants of AHA<sup>-</sup>. Both formyl hydrogen-deuterium substitution (Figure 2c) and carbon-13 substitution (Supporting Information, Figure S1) were found to result in only minor changes in the spectra obtained in helium nanodroplets. In contrast, the spectrum of the fully deuterated dimer (ADA<sup>-</sup>), in which the shared proton has been exchanged for a deuteron, differs significantly from that of the unlabeled species, as shown in Figure 2 d. Most notably, a shift of the most intense spectral band from 605 cm<sup>-1</sup> ( $b_1$ , Figure 2b) to 433 cm<sup>-1</sup> ( $d_1$ , Figure 2d) is observed in the helium nanodroplet spectra. The spectrum of ADA<sup>-</sup> measured with Ar tagging in the currently accessible photon energy range  $(>600 \text{ cm}^{-1}, \text{ Figure 2e})$ agrees well with that obtained by the helium nanodroplet method (Supporting Information, Table S1).

Angewandte

Chemie

The substantial H/D isotopic shift in the energy of band  $b_1$  provides strong evidence that this feature is associated with displacement of the shared proton. In studies of other anionic proton-bound dimers, such intense spectral lines have been traced to the fundamental transition associated with proton motion parallel to the shared proton axis,  $v_{\parallel}$ .<sup>[17,20a,22]</sup> Interestingly, however, calculation of the frequency ratio between band  $b_1$  of AHA<sup>-</sup> and band  $d_1$  of ADA<sup>-</sup> yields a value of 1.40, which is inconsistent with the significantly larger frequency shift expected for an anharmonic potential well (for example, a ratio of about 1.5 for deuterium substitution in a quartic potential).<sup>[23]</sup> Thus, the vibrational mode associated with this band cannot be confidently assigned solely from the changes in band energies upon isotopic substitution.

To gauge the degree of coupling between the displacements of the shared proton and the deformations of the flanking HCOO- ions, potential energy surfaces (PESs) of structure 1 in a normal coordinate basis were constructed and solved for their eigenstates and eigenvectors utilizing the discrete variable representation (DVR) method,[20b,24] as detailed in the Supporting Information. In the harmonic approximation, two intense bands below 1200 cm<sup>-1</sup> are predicted by MP2 calculations for AHA<sup>-</sup> (f<sub>1</sub> and f<sub>2</sub>, Figure 3b), corresponding to the fundamental of  $v_{\parallel}$  and out-ofphase carboxylate deformation,  $\delta(OCO)_{OOP}$  respectively. However, utilizing one-dimensional (1D) PESs in the corresponding normal coordinate, the first excited states of  $\nu_{\scriptscriptstyle \|}$  and  $\delta(OCO)_{OOP}$  are calculated to be nearly exactly isoenergetic, circa 899 cm<sup>-1</sup> above the vibrational ground state (Supporting Information, Table S2). Since both modes belong to the B symmetry species of the C<sub>2</sub> point group, the fundamental excitations couple strongly to yield mixed-character vibrational modes that can be represented as a linear combination of the contributing normal mode excited states on a twodimensional (2D) PES (Supporting Information, Figure S4). We find that the low-energy vibrational level (657  $cm^{-1}$  above the ground state) is composed of 42 %  $\delta(OCO)_{OOP}$  and 52 %  $\nu_{\parallel}$  first excited states, and the high-energy vibrational level (1213  $\text{cm}^{-1}$  above the ground state) is composed of 52%  $\delta(OCO)_{OOP}$  and 40% v<sub>I</sub> first excited states. Higher excited states of  $\delta(OCO)_{OOP}$  and  $v_{\parallel}$  contribute less than 5% to the coupled states (Supporting Information, Table S3). These fractional contributions can also be applied to the Cartesian displacement vectors associated with the normal modes to evaluate the molecular motion for the coupled modes. Whereas normal modes  $\nu_{_{\parallel}}$  and  $\delta(OCO)_{OOP}$  both involve large-amplitude displacement of the shared proton along the O-H-O axis (Supporting Information, Figure S5a,b), mode coupling largely confines shared proton motion to the highenergy coupled mode (Supporting Information, Figure S5c), and carboxylate deformation is observed in the low-energy mode (Supporting Information, Figure S5d). Animations of the coupled modes are available as Supporting Information.

In ADA<sup>-</sup>, the exchange of the shared proton for a deuteron leads to an energetic separation of the first excited states of  $v_{\parallel}$  and  $\delta(OCO)_{OOP}$  which are calculated from 1D PESs to lie 595 cm<sup>-1</sup> and 812 cm<sup>-1</sup> above the ground state, respectively (Supporting Information, Table S2). As a result, the vibrational levels remain largely decoupled in 2D PES calculations, with a low-energy level (544 cm<sup>-1</sup> above the ground state) composed of 15%  $\delta(OCO)_{OOP}$  and 82%  $v_{\parallel}$  and a high-energy level (891 cm<sup>-1</sup> above the ground state) composed of 83%  $\delta(OCO)_{OOP}$  and 15%  $v_{\parallel}$  (Supporting Information, Table S3). This result clearly demonstrates an isotope-dependent coupling of vibrational normal modes, providing a physical explanation for the smaller-thanexpected frequency shift of band  $b_1$  upon deuterium substitution.

Although strong coupling between  $v_{\parallel}$  and  $\delta(OCO)_{OOP}$  is observed in the 2D PES for AHA-, the predicted frequencies and intensities are insufficient to explain the bands observed experimentally (Supporting Information, Table S2). To better account for coupling of proton motion to the interformate O-O distance, a four-dimensional (4D) PES was constructed for both AHA<sup>-</sup> and ADA<sup>-</sup>, adding the O-H-O symmetric stretch,  $v_s$ (O-H-O), and in-phase carboxylate deformation,  $\delta(OCO)_{IP}$  to the normal coordinate basis. Improved agreement in the predicted frequencies and intensities is observed utilizing this PES, with intense bands predicted for AHA<sup>-</sup> at 619, 944, and 1103  $\text{cm}^{-1}$  (bands g1, g2, and g3; Figure 3c) and for ADA<sup>-</sup> at 466, 786, and 918 cm<sup>-1</sup> (bands  $j_1$ ,  $j_2$ , and  $j_3$ , Figure 3g). A decomposition of the four-dimensional eigenvectors as a linear combination of normal mode eigenvectors shows significant contributions from states involving simultaneous excitation of multiple normal modes (that is, combination bands; Supporting Information, Tables S5, S6). We anticipate that more comprehensive multidimensional potential calculations, for example those that include formate torsional modes, will yield further insight into the origin of low-energy bands in the spectrum of AHA<sup>-</sup> and ADA<sup>-</sup>.

In summary, we have characterized the ground-state structure of the proton-bound formate dimer,  $HCOO^{-}\cdot H^{+,-}OOCH$ , utilizing vibrational spectroscopy of ions trapped in helium nanodroplets and photodissociation of

Ar-complexed ions. Analysis of the CO<sub>2</sub><sup>-</sup> normal modes supports a symmetrical structure for the dimer with an equally shared proton. Preliminary multidimensional potential calculations reveal strong coupling between the vibrational normal modes and also show the strong isotope dependence of the mode coupling. It is intriguing to consider how the carboxylate substituent moiety may influence the structure and dynamics of proton-bound dimers, as the ground-state geometry and coupling of normal modes are likely highly sensitive to the identity of this functional group. Furthermore, knowing that the proton is localized in the intramolecular proton-bound dicarboxylate motif of deprotonated dodecanedioic acid,<sup>[8]</sup> the question arises as to which structural constraints dictate the transition from shared to localized proton. These queries present appealing candidates for future studies to obtain a more comprehensive picture of the structure and dynamics of the prevalent proton-bound dicarboxylate motif.

#### Acknowledgements

The authors thank Christopher Leavitt for acquisition of the Ar-tagging data and the FHI FEL staff, particularly Sandy Gewinner and Wieland Schöllkopf, for laser operation. D.A.T. gratefully acknowledges the support of the Alexander von Humboldt Foundation. M.A.J. thanks the National Science Foundation under grant CHE-1465100.

### **Conflict of interest**

The authors declare no conflict of interest.

Keywords: carboxylates  $\cdot$  cold-ion spectroscopy  $\cdot$  coupling  $\cdot$  helium nanodroplets  $\cdot$  shared protons

How to cite: Angew. Chem. Int. Ed. 2018, 57, 10615–10619 Angew. Chem. 2018, 130, 10775–10779

- a) G. C. Pimentel, J. Chem. Phys. **1951**, 19, 446-448; b) K. Kawaguchi, E. Hirota, J. Chem. Phys. **1986**, 84, 2953-2960; c) K. Kawaguchi, E. Hirota, J. Chem. Phys. **1987**, 87, 6838-6841; d) G. Pérez-Hernández, J. González-Vázquez, L. González, J. Phys. Chem. A **2012**, 116, 11361-11369.
- [2] H. Rodríguez, G. Gurau, J. D. Holbrey, R. D. Rogers, *Chem. Commun.* 2011, 47, 3222–3224.
- [3] G. Gurau, H. Rodríguez, S. P. Kelley, P. Janiczek, R. S. Kalb, R. D. Rogers, *Angew. Chem. Int. Ed.* 2011, 50, 12024–12026; *Angew. Chem.* 2011, 123, 12230–12232.
- [4] J. Lin, E. Pozharski, M. A. Wilson, *Biochemistry* 2017, 56, 391– 402.
- [5] a) A. I. González Flórez, E. Mucha, D.-S. Ahn, S. Gewinner, W. Schöllkopf, K. Pagel, G. von Helden, *Angew. Chem. Int. Ed.* **2016**, *55*, 3295–3299; *Angew. Chem.* **2016**, *128*, 3356–3360; b) E. Mucha, A. I. González Flórez, M. Marianski, D. A. Thomas, W. Hoffmann, W. B. Struwe, H. S. Hahm, S. Gewinner, W. Schöllkopf, P. H. Seeberger, G. von Helden, K. Pagel, *Angew. Chem. Int. Ed.* **2017**, *56*, 11248–11251; *Angew. Chem.* **2017**, *129*, 11400–11404.
- [6] A. B. Wolk, C. M. Leavitt, E. Garand, M. A. Johnson, Acc. Chem. Res. 2014, 47, 202–210.

- [7] a) K. R. Asmis, Y. Yang, G. Santambrogio, M. Brümmer, J. R. Roscioli, L. R. McCunn, M. A. Johnson, O. Kühn, Angew. Chem. Int. Ed. 2007, 46, 8691-8694; Angew. Chem. 2007, 119, 8846-8849; b) J. R. Roscioli, L. R. McCunn, M. A. Johnson, Science 2007, 316, 249-254; c) C. J. Johnson, A. B. Wolk, J. A. Fournier, E. N. Sullivan, G. H. Weddle, M. A. Johnson, J. Chem. Phys. 2014, 140, 221101; d) J. A. Fournier, C. T. Wolke, M. A. Johnson, T. T. Odbadrakh, K. D. Jordan, S. M. Kathmann, S. S. Xantheas, J. Phys. Chem. A 2015, 119, 9425-9440; e) C. T. Wolke, J. A. Fournier, L. C. Dzugan, M. R. Fagiani, T. T. Odbadrakh, H. Knorke, K. D. Jordan, A. B. McCoy, K. R. Asmis, M. A. Johnson, Science 2016, 354, 1131-1135; f) Y. Yang, O. Kühn, G. Santambrogio, D. J. Goebbert, K. R. Asmis, J. Chem. Phys. 2008, 129, 224302; g) D. T. Moore, J. Oomens, L. van der Meer, G. von Helden, G. Meijer, J. Valle, A. G. Marshall, J. R. Eyler, ChemPhysChem 2004, 5, 740-743.
- [8] M. Z. Kamrath, R. A. Relph, T. L. Guasco, C. M. Leavitt, M. A. Johnson, *Int. J. Mass Spectrom.* 2011, 300, 91–98.
- [9] A. Halkier, T. Helgaker, P. Jørgensen, W. Klopper, J. Olsen, *Chem. Phys. Lett.* **1999**, 302, 437–446.
- [10] R. A. Kendall, T. H. Dunning, Jr., R. J. Harrison, J. Chem. Phys. 1992, 96, 6796-6806.
- [11] C. Møller, M. S. Plesset, Phys. Rev. 1934, 46, 618-622.
- [12] a) K. Raghavachari, G. W. Trucks, J. A. Pople, M. Head-Gordon, *Chem. Phys. Lett.* **1989**, *157*, 479–483; b) R. J. Bartlett, J. D. Watts, S. A. Kucharski, J. Noga, *Chem. Phys. Lett.* **1990**, *165*, 513–522.
- [13] M. O. Sinnokrot, E. F. Valeev, C. D. Sherrill, J. Am. Chem. Soc. 2002, 124, 10887–10893.

[14] W. H. Robertson, J. A. Kelley, M. A. Johnson, J. Chem. Phys. 2000, 113, 7879-7884.

Angewandte

Chemie

- [15] H. K. Gerardi, A. F. DeBlase, X. Su, K. D. Jordan, A. B. McCoy, M. A. Johnson, J. Phys. Chem. Lett. 2011, 2, 2437–2441.
- [16] I. D. Reva, A. M. Plokhotnichenko, E. D. Radchenko, G. G. Sheina, Y. P. Blagoi, *Spectrochim. Acta Part A* 1994, 50, 1107– 1111.
- [17] E. G. Diken, J. M. Headrick, J. R. Roscioli, J. C. Bopp, M. A. Johnson, A. B. McCoy, J. Phys. Chem. A 2005, 109, 1487–1490.
- [18] a) F. Filsinger, D.-S. Ahn, G. Meijer, G. von Helden, *Phys. Chem. Chem. Phys.* 2012, *14*, 13370–13377; b) A. I. González Flórez, D.-S. Ahn, S. Gewinner, W. Schöllkopf, G. von Helden, *Phys. Chem. Chem. Phys.* 2015, *17*, 21902–21911.
- [19] C. T. Wolke, A. F. DeBlase, C. M. Leavitt, A. B. McCoy, M. A. Johnson, J. Phys. Chem. A 2015, 119, 13018-13024.
- [20] a) N. Heine, T. I. Yacovitch, F. Schubert, C. Brieger, D. M. Neumark, K. R. Asmis, J. Phys. Chem. A 2014, 118, 7613-7622;
   b) J. A. Tan, J.-L. Kuo, J. Phys. Chem. A 2015, 119, 11320-11328.
- [21] L. R. McCunn, J. R. Roscioli, M. A. Johnson, A. B. McCoy, J. Phys. Chem. B 2008, 112, 321–327.
- [22] N. Heine, K. R. Asmis, Int. Rev. Phys. Chem. 2014, 34, 1-34.
- [23] R. P. Bell, Proc. R. Soc. London Ser. A 1945, 183, 328-337.
- [24] a) D. T. Colbert, W. H. Miller, J. Chem. Phys. 1992, 96, 1982–1991; b) J. C. Light, T. Carrington, Advances in Chemical Physics, Wiley, Hoboken, 2007, pp. 263–310.

Manuscript received: May 10, 2018 Accepted manuscript online: June 19, 2018 Version of record online: July 12, 2018